# Self-Calibration in Radio Astronomy

by Subhashis Roy

## Insufficiency of Standard Calibration

$$V'_{ij}(t, u, v) = g_i(t).g_j^*(t).V_{ij}(u, v)$$

g(t) varies with time during Target source observations.

Main problem from Ionospheric phase variation.

Causes point source to smear out and low level fake structures in a map.

Gain error:  $\sigma_G^2 = \frac{\sigma_V^2}{(N-3)S^2} \ (\text{N is no. of antennas}) \, .$ 

### Self-calibration

N complex gain errors corrupt  $\left(\frac{N(N-1)}{2}\right)$  visibilities.

Allowing N Gains to be variable, provide  $\left(\frac{N(N-1)}{2}\right)-N$  constraints.

#### Self-calibration involves minimising

$$S = \sum_{k} \sum_{i,j,i \neq j} |V_{ij}(t_k) - g_i(t_k)g_j^*(t_k).V_{ij}^M(t_k)|^2$$

or 
$$S = \sum_k \sum_{i,j,i\neq j} |V_{ij} \rangle |^2 |X_{ij}(t_k) - g_i(t_k)g_j^*(t_k)|^2 \\ \longrightarrow (1)$$
 where  $X_{ij}(t_k) = \frac{V_{ij}(t_k)}{V_{ij}^M(t_k)}$ .

#### Steps:

- 1. Make an initial model of the source using preliminary Cal. and acceptable source structures.
- 2. Solve for complex gains using Eq.(1).
- 3. Find the corrected visibility  $V_{\rm ij,corr}(t)=rac{V_{\rm ij}(t)}{g_i(t)g_j^*(t)}$
- 5. Form a new model from the image made from the corrected data.
- 6. Go to 2, unless satisfied.

[Aver aging time]

#### How does it work

Consider

$$I(l,m) = \frac{1}{M} \sum_{k=1}^{M} V(u_k,v_k) e^{2\pi i (u_k l + v_k m)}$$
 in 1-D.

$$I(l) = \frac{1}{M} \sum_{k=1}^{M} V(u_k) e^{2\pi i (u_k l)}$$

Assume N-1 antennas to have 0 phases. 1 to have **1 Rad** as phase.

Real source of unit strength at l=0. Recall

$$V'_{ij}(t, u, v) = g_i(t).g_j^*(t).V_{ij}(u, v)$$

$$I(l) = \frac{1}{M} \left( \sum_{k=1}^{M-(N-1)} e^{2\pi i(u_k l)} + \sum_{M-N}^{M} e^{i} \cdot e^{2\pi i(u_k l)} \right)$$